

# $K_{3,3}$ -free Intersection Graphs of Finite Groups\*

Selçuk Kayacan<sup>†</sup>

Department of Mathematics, Istanbul Technical University, 34469 Maslak, Istanbul,  
Turkey.

*e-mail:* [skayacan@itu.edu.tr](mailto:skayacan@itu.edu.tr)

## Abstract

The intersection graph of a group  $G$  is an undirected graph without loops and multiple edges defined as follows: the vertex set is the set of all proper non-trivial subgroups of  $G$ , and there is an edge between two distinct vertices  $H$  and  $K$  if and only if  $H \cap K \neq 1$  where  $1$  denotes the trivial subgroup of  $G$ . In this paper we classify all finite groups whose intersection graphs are  $K_{3,3}$ -free.

2010 *Mathematics Subject Classification*. Primary: 20D99; Secondary: 20D15, 20D25, 05C25.

Keywords: Finite groups; subgroup; intersection graph

## Introduction

Let  $\bar{\Gamma}$  and  $\Gamma$  be two graphs. We say that  $\bar{\Gamma}$  is  $\Gamma$ -free if there is no subgraph of  $\bar{\Gamma}$  which is isomorphic to  $\Gamma$ , i.e.  $\bar{\Gamma}$  does not contain  $\Gamma$  as a subgraph. Let  $G$  be a group. We denote by  $\Gamma(G)$  the intersection graph (of subgroups) of  $G$ . For simplicity we say that  $G$  is  $\Gamma$ -free whenever its intersection graph is  $\Gamma$ -free.

In a recent work [9], Rajkumar and Devi classified finite groups whose intersection graphs does not contain one of  $K_5$ ,  $K_4$ ,  $C_5$ ,  $C_4$ ,  $P_4$ ,  $P_3$ ,  $P_2$ ,  $K_{1,3}$ ,  $K_{2,3}$  or  $K_{1,4}$  as a subgraph. Here we present the classification of finite  $K_{3,3}$ -free groups. Our main result is

**Theorem.** *A finite group is  $K_{3,3}$ -free if and only if it is isomorphic to the one of the following groups:*

1.  $\mathbb{Z}_{pqr}$ ,  $\mathbb{Z}_{p^2q}$ ,  $\mathbb{Z}_{pq}$ ,  $\mathbb{Z}_{p^i}$ , where  $p, q, r$  are distinct primes and  $0 \leq i \leq 6$ .
2.  $\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_p \times \mathbb{Z}_p$ ,  $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_p$  ( $p \neq 2$ ), where  $p$  is a prime.

---

\*I would like to thank to the anonymous referee for the meticulous reading and the shorter arguments he/she provides which greatly enhances the exposition. This work is completed while I was a visitor in Graz University of Technology. By this occasion I would like to thank to Math C Department of TUGraz for their hospitality.

<sup>†</sup>This work is supported by the TÜBİTAK 2214/A Study Abroad Fellowship Program: 1059B141401085.

3. The dihedral group  $D_8$  of order 8, the quaternion group  $Q_8$  of order 8.
4. The semidirect products  $\mathbb{Z}_q \rtimes \mathbb{Z}_{p^2}$  with  $p^2 \mid q-1$ ,  $(\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_q$  with  $q \mid p+1$ , where  $p, q$  are distinct primes.
5. The semidirect product  $(\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_{q^2}$  with  $q^2 \mid p+1$ , where  $p, q$  are distinct primes.
6. The semidirect product  $\mathbb{Z}_r \rtimes \mathbb{Z}_{pq}$  with  $pq \mid r-1$ , where  $p, q, r$  are distinct primes.
7. The semidirect product  $\mathbb{Z}_p \rtimes \mathbb{Z}_q$  with  $q \mid p-1$ , where  $p > q$  are distinct primes.
8.  $\mathbb{Z}_{p^3} \times \mathbb{Z}_q$ ,  $\mathbb{Z}_9 \times \mathbb{Z}_3$ ,  $(\mathbb{Z}_3 \times \mathbb{Z}_3) \rtimes \mathbb{Z}_3$ ,  $\mathbb{Z}_9 \rtimes \mathbb{Z}_3$ ,  $\mathbb{Z}_3 \rtimes \mathbb{Z}_4$ , and  $D_{18}$ , where  $p, q$  are distinct primes.
9. The semidirect product  $\mathbb{Z}_q \rtimes \mathbb{Z}_{p^3}$  with  $p^3 \mid q-1$ , where  $p, q$  are distinct primes.

In [8], Kayacan and Yaraneri classified all finite groups whose intersection graphs are planar (planar groups in short). By Kuratowski's theorem a graph is planar if and only if it does not contain a subgraph that is a subdivision of  $K_5$  or  $K_{3,3}$ . Therefore the above list contains all finite groups with planar intersection graphs. Actually, the first seven items lists the planar groups (except  $\mathbb{Z}_{p^6}$ ) and the last two items presents the  $K_{3,3}$ -free groups containing a  $K_5$  as a subgraph in their intersection graphs. Our methods are elementary. Occasionally, we used some improved versions of the arguments that are presented in [8]. Notation is standard and should be clear to the reader. Below we present some of the results we used in the text.

*Remarks.*

1. (see [10, p. 30]) (Product Formula)  $|XY||X \cap Y| = |X||Y|$  for any two subgroups  $X$  and  $Y$  of a finite group.
2. (see [5, p. 30]) (Sylow Theorems)
  - (a) Let  $G$  be a group of order  $p^n$  and  $k < n$ . Then the number of subgroups of order  $p^k$  in  $G$  is  $\equiv 1 \pmod{p}$ .
  - (b) Let  $G$  be a group of order  $p^n s$ ,  $p \nmid s$ ,  $k \leq n$ . Then the number of subgroups of order  $p^k$  in  $G$  is  $\equiv 1 \pmod{p}$ .
3. (see [10, p. 81]) Let  $P$  be a Sylow  $p$ -subgroup of a finite group  $G$ . Then,  $N_G(H) = H$  for any subgroup  $H$  with  $N_G(P) \leq H \leq G$ .
4. (see [7, p. 24]) In a finite solvable group  $G$ , the factors of every chief series are elementary abelian of prime power order. In particular, every minimal normal subgroup of  $G$  is elementary abelian.
5. (see [7, p. 231]) If  $G$  is a finite solvable group, then any  $\pi$ -subgroup is contained in a Hall  $\pi$ -subgroup. Moreover, any two Hall  $\pi$ -subgroup are conjugate.
6. (see [10, p. 197]) Let  $G$  be a finite group and  $p$  be the smallest prime divisor of  $|G|$ . If a Sylow  $p$ -subgroup  $P$  of  $G$  is cyclic, then there is a normal subgroup  $N$  of  $G$  such that  $P \cap N = 1$  and  $G = PN$ .

## Nilpotent groups

Obviously, if  $G$  is  $\Gamma$ -free, then any subgroup of  $G$  is also  $\Gamma$ -free. With this simple remark and by using the fundamental theorem of finite abelian groups, it is not difficult to determine  $K_{3,3}$ -free abelian groups.

**Lemma 1.** *A finite abelian group is  $K_{3,3}$ -free if and only if it is isomorphic to one of the following groups*

$$\begin{aligned} \mathbb{Z}_{p^i} \ (0 \leq i \leq 6), \quad \mathbb{Z}_{p^3} \times \mathbb{Z}_q, \quad \mathbb{Z}_{p^2} \times \mathbb{Z}_q, \quad \mathbb{Z}_p \times \mathbb{Z}_q, \quad \mathbb{Z}_9 \times \mathbb{Z}_3, \quad \mathbb{Z}_4 \times \mathbb{Z}_2, \\ \mathbb{Z}_p \times \mathbb{Z}_p, \quad \mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r, \quad \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_p \ (p \neq 2) \end{aligned}$$

where  $p$ ,  $q$ , and  $r$  are distinct primes.

*Proof.* Let  $G$  be a finite abelian group and  $p$ ,  $q$ , and  $r$  be distinct prime numbers.

*Case I:*  $G$  is a cyclic group. Then there is exactly one subgroup of  $G$  of order  $n$  for each divisor  $n$  of  $|G|$ . Observe that  $\Gamma(G)$  is  $K_{3,3}$ -free if  $G$  is of order  $p^i$  ( $0 \leq i \leq 6$ ),  $p^2q$ , or  $pq$ ; as the number of proper non-trivial subgroups of  $G$  would be less than six in such cases.

*Case I (a):*  $|G| = p^i$  ( $i > 6$ ). Then  $\Gamma(G)$  contains a  $K_6$  and cannot be  $K_{3,3}$ -free.

*Case I (b):*  $G \cong \mathbb{Z}_{p^3q}$ . Let  $a$  and  $b$  be two elements of  $G$  of order  $p^3$  and of order  $q$  respectively. There are exactly six proper non-trivial subgroups of  $G$  in this case and five of them, namely  $\langle a^{p^2} \rangle$ ,  $\langle a^p \rangle$ ,  $\langle a \rangle$ ,  $\langle a^{p^2}, b \rangle$ , and  $\langle a^p, b \rangle$  form a complete graph in  $\Gamma(G)$  as all of them contain  $\langle a^{p^2} \rangle$ . The remaining vertex  $\langle b \rangle$  has degree two in the intersection graph and, therefore,  $\Gamma(G)$  is  $K_{3,3}$ -free. Moreover, since  $\Gamma(G)$  contains a  $K_5$ , it cannot be a proper subgroup of a  $K_{3,3}$ -free group. (Notice that in a larger group  $G$  becomes a vertex and so vertices containing  $\langle a^{p^2} \rangle$  form a subgraph containing a  $K_6$ .)

*Case I (c):*  $G \cong \mathbb{Z}_{p^2q^2}$ . Let  $a$  and  $b$  be two elements of  $G$  of order  $p^2$  and of order  $q^2$  respectively. As in the previous case we have five subgroups forming a  $K_5$ , namely  $\langle a^p \rangle$ ,  $\langle a \rangle$ ,  $\langle a^p, b^q \rangle$ ,  $\langle a, b^q \rangle$ , and  $\langle a^p, b \rangle$ . However, unlike the previous case, the subgroup  $\langle b^q \rangle$  is linked by an edge with those last three subgroups forming the  $K_5$ . Therefore,  $\Gamma(G)$  is not  $K_{3,3}$ -free in this case.

*Case I (d):*  $G \cong \mathbb{Z}_{pqr}$ . Let  $a, b$ , and  $c$  be three elements of orders  $p, q$ , and  $r$  respectively. The vertices of  $\Gamma(G)$  are  $\langle a \rangle$ ,  $\langle b \rangle$ ,  $\langle c \rangle$ ,  $\langle a, b \rangle$ ,  $\langle a, c \rangle$ , and  $\langle b, c \rangle$ . As the valency of  $\langle a \rangle$  is two,  $\Gamma(G)$  is  $K_{3,3}$ -free in this case. However, if  $G$  is a proper subgroup of a larger group, then maximal subgroups of  $G$  together with  $\langle a \rangle$ ,  $\langle b \rangle$ , and  $G$  form a subgraph containing a  $K_{3,3}$ .

To sum up, the only possible values for the order of a  $K_{3,3}$ -free cyclic group are

$$p^i \ (0 \leq i \leq 6), \quad p^3q, \quad p^2q, \quad pqr, \quad pq.$$

*Case II:*  $G$  is not a cyclic group. Let us make a useful observation. If  $G$  is a  $K_{3,3}$ -free abelian group of order  $n$ , then  $n$  must be one of the above values. This is because for any pair of subgroups  $A < B$  of  $\mathbb{Z}_n$ , there are corresponding subgroups  $H < K$  of  $G$  such that  $|A| = |H|$  and  $|B| = |K|$ .

*Case II (a):*  $G \cong \mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p$ . Observe that maximal subgroups of  $G$  are dimension 2 subspaces of  $G$  considering it as a vector space over  $\mathbb{F}_p$ . Then, by a counting argument the number of maximal subgroups of  $G$  is  $[(p^3 - 1)(p^3 - p)]/[(p^2 - 1)(p^2 - p)] = p^2 + p + 1$ . However, by the product formula (see Remark 1), any pair of maximal subgroups intersects non-trivially; and hence, they form a complete graph in  $\Gamma(G)$ . Thus, any group containing an elementary abelian subgroup of rank three cannot be  $K_{3,3}$ -free.




Figure 1

*Case II (b):*  $G \cong \mathbb{Z}_{p^2} \times \mathbb{Z}_p$ . Let  $a$  and  $b$  be two generator elements for  $G$  of order  $p^2$  and of order  $p$  respectively. Then, subgroups  $\langle a^p \rangle, \langle a^p, b \rangle, \langle a \rangle, \langle ab \rangle, \dots, \langle ab^{p-1} \rangle$  form a  $K_{p+2}$  in  $\Gamma(G)$ . Hence, the only possible values of  $p$  are 2 and 3, if  $G$  is  $K_{3,3}$ -free. Actually,  $G$  is  $K_{3,3}$ -free for those primes. Intersection graph of  $\mathbb{Z}_9 \times \mathbb{Z}_3$  is depicted in Figure 1a.

*Case II (c):*  $G$  is an abelian  $p$ -group which is not considered in Cases II (a) and (b). If  $G \cong \mathbb{Z}_p \times \mathbb{Z}_p$  then  $\Gamma(G)$  is  $K_{3,3}$ -free, since the intersection graph of a group of order  $p^2$  consists of isolated vertices. Otherwise,  $G$  contains a subgroup  $H$  properly which is isomorphic to  $\mathbb{Z}_{p^2} \times \mathbb{Z}_p$  with  $p \in \{2, 3\}$  by the previous cases. Let  $a$  and  $b$  be two generator elements for  $H$  of order  $p^2$  and of order  $p$  respectively. Since  $H$  is a proper subgroup of  $G$ , there exists an element  $c \in G$  that does not lie in  $H$ . Now, if  $c^p \in \langle a \rangle$ , then  $\langle a^p \rangle, \langle a^p, b \rangle, \langle a \rangle, \langle ab \rangle, \langle c \rangle$ , and  $H$  form a  $K_6$  in  $\Gamma(G)$ . And if  $c^p \notin \langle a \rangle$ , then  $\langle a^p \rangle, \langle a^p, b \rangle, \langle a \rangle, \langle ab \rangle, \langle a^p, c \rangle$ , and  $H$  form a  $K_6$  in  $\Gamma(G)$ .

*Case II (d):*  $|G| = p^3q$ . Since  $G$  is a non-cyclic abelian group, either  $G \cong (\mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p) \times \mathbb{Z}_q$  or  $G \cong (\mathbb{Z}_{p^2} \times \mathbb{Z}_p) \times \mathbb{Z}_q$ . However, the first case cannot occur if  $G$  is  $K_{3,3}$ -free in virtue of Case II (a); and in the latter case  $\langle a^p \rangle, \langle a^p, b \rangle, \langle a \rangle, \langle ab \rangle, \langle a, b \rangle$ , and  $\langle a^p, c \rangle$  form a  $K_6$  in the intersection graph where  $a, b$ , and  $c$  are some generators of  $G$  of orders  $p^2, p$ , and  $q$  respectively.

*Case II (e):*  $G \cong (\mathbb{Z}_p \times \mathbb{Z}_p) \times \mathbb{Z}_q$ . Let  $a$  and  $b$  be two elements of  $G$  generating a subgroup of order  $p^2$ , and let  $c$  be an element of  $G$  of order  $q$ . As in Case II (b), subgroups  $\langle c \rangle, \langle a, c \rangle, \langle ab, c \rangle, \dots, \langle ab^{p-1}, c \rangle$ , and  $\langle b, c \rangle$  form a  $K_{p+2}$  in  $\Gamma(G)$ , and therefore, if  $G$  is  $K_{3,3}$ -free then either  $p = 2$  or  $p = 3$ . If  $p = 2$ , then  $G$  is  $K_{3,3}$ -free and its intersection graph presented in Figure 1b. However, if  $p = 3$ , then subgroups  $\langle c \rangle, \langle a, c \rangle, \langle ab, c \rangle, \langle ab^2, c \rangle, \langle b, c \rangle$  together with  $\langle a, b \rangle$  form a subgraph containing  $K_{3,3}$ .

As abelian groups of order  $pqr$  and of order  $pq$  are necessarily cyclic, the proof is completed.  $\square$

Let  $G$  be a non-abelian  $p$ -group of order  $p^\alpha$  ( $\alpha > 2$ ) which is  $K_{3,3}$ -free. Then, as the quotient of  $G$  by the Frattini subgroup  $\Phi(G)$  (i.e. the intersection of all maximal subgroups) is elementary abelian,  $\Phi(G)$  is a non-trivial subgroup of  $G$ . That is, there are subgroups  $K, L$  of  $G$  such that  $|K| = p$ ,  $|L| = p^2$ , and  $K$  is contained by every maximal subgroup of  $G$  as well as by  $L$ . If  $p > 3$ , by Remark 2 there are at least 6 maximal subgroups each containing a common subgroup, hence  $\Gamma(G)$  contains a  $K_7$  which is a contradiction. (Notice that for a graph being  $K_{3,3}$ -free is a more stringent condition than being  $K_6$ -free.) Thus  $G$  is either a 2-group or a 3-group. Also, the exponent  $\alpha = 3$ . To see this, suppose that  $\alpha = 4$  and consider the case  $p = 3$ . Then there are at least four




Figure 2

maximal subgroups of  $G$  of order  $p^3$  and together with  $K$  and  $L$ , they form a  $K_6$  in the intersection graph. If  $\alpha = 4$ ,  $p = 2$  and  $|\Phi(G)| = p$ , then  $G/\Phi(G)$  is elementary abelian of rank 3 which is not listed in Lemma 1. Suppose that  $|\Phi(G)| = p^2$ . If  $\Phi(G)$  is cyclic, then the subgroup  $Z \leq \Phi(G)$  of order  $p$  is in the center of  $G$ . Since the intersection graph of the quaternion group of order 16 is  $K_9$ , we may further assume there are more than one minimal subgroups of  $G$ . Let  $K$  be another minimal subgroup of  $G$ , then three maximal subgroups together with  $Z$ ,  $\Phi(G)$ , and  $ZK$  form a  $K_6$  in  $\Gamma(G)$ . If  $\Phi(G)$  is not cyclic, then we may take three maximal subgroups and the three subgroups of  $\Phi(G)$  to form a  $K_{3,3}$  in the intersection graph. Finally,  $|\Phi(G)| = p^3$  implies  $G$  is cyclic which is a contradiction. There are two non-abelian groups of order 8, namely the dihedral group  $D_8$  and the quaternion group  $Q_8$ ; and also there are two non-abelian groups of order 27, namely  $(\mathbb{Z}_3 \times \mathbb{Z}_3) \rtimes \mathbb{Z}_3$  and  $\mathbb{Z}_9 \rtimes \mathbb{Z}_3$ . It can be verified that these groups are  $K_{3,3}$ -free (see Figure 2 and Figure 3). Thus, we almost proved that

**Lemma 2.** *A finite non-abelian nilpotent group is  $K_{3,3}$ -free if and only if it is isomorphic to one of the following groups*

$$D_8, \quad Q_8, \quad (\mathbb{Z}_3 \times \mathbb{Z}_3) \rtimes \mathbb{Z}_3, \quad \mathbb{Z}_9 \rtimes \mathbb{Z}_3.$$

*Proof.* Since a nilpotent group  $G$  is the direct product of its Sylow subgroups, at least one of them must be non-abelian. However,  $(\mathbb{Z}_3 \times \mathbb{Z}_3) \rtimes \mathbb{Z}_3$  and  $\mathbb{Z}_9 \rtimes \mathbb{Z}_3$  both contains a  $K_5$  in their intersections graphs, therefore cannot be a proper subgroup of  $G$ . Also, if  $G$  contains  $D_8$  properly, then the three maximal subgroups together with  $D_8$  and  $\Phi(D_8)$  form a  $K_5$  in  $\Gamma(G)$ . If we take another minimal subgroup  $K$  which is not a subgroup of  $D_8$ , then the subgroup  $\Phi(D_8)K$  would be a sixth vertex which is connected by an edge with each vertices of  $K_5$ . Same argument is valid also for  $Q_8$ .  $\square$

## Solvable groups

Let  $G$  be a finite non-nilpotent solvable group. Following [8, Sec. 4], we may reduce the number of cases substantially with regard to the orders of the groups. Let  $N$  be a minimal subgroup of  $G$ . By Remark 4,  $N$  is an elementary abelian group and as a subgroup of  $G$  it is  $K_{3,3}$ -free. Moreover,  $N$  is either of rank 1 or rank 2 in virtue of Lemma 1. It is well-known that there is a correspondence between the subgroups of  $G$  containing  $N$  and the subgroups of  $G/N$ . Now we make a very useful observation

- the rank of  $N$  is 1  $\implies$  # subgroups of  $G/N$  is at most 6;
- the rank of  $N$  is 2  $\implies$  # subgroups of  $G/N$  is at most 3.




Figure 3: Vertices labelled with  $\Phi$  represents Frattini subgroups

As a consequence of Sylow and Hall Theorems, the only possible values of  $|G/N|$  are  $p^i$  ( $0 \leq i \leq 5$ ),  $pq$ , and  $p^2q$  where  $p$  and  $q$  distinct prime numbers. Moreover, if  $N$  is of rank 2, then  $G/N$  is isomorphic to a cyclic group of prime or prime squared order. Therefore, the only possible cases for the order of  $G$  are

$$p^5q, \quad p^4q, \quad p^2qr; \quad p^3q, \quad p^2q, \quad p^2q^2, \quad pqr, \quad pq.$$

In [9], the  $K_{2,3}$ -free groups are determined as a sublist of  $K_5$ -free groups. Our preceding discussion made it apparent that if  $G$  is a  $K_{2,3}$ -free group and  $N \triangleleft G$  is elementary abelian of rank 2, then either  $|G : N| = 1$  or  $p$  where  $p$  denotes a prime number.

Returning to the possible orders of the non-nilpotent solvable  $K_{3,3}$ -free groups, we can still eliminate some of the cases by *ad hoc* arguments.

**Lemma 3.** *There are no finite non-nilpotent solvable group which is  $K_{3,3}$ -free and of order*

$$p^5q, \quad p^4q, \quad \text{or} \quad p^2qr$$

where  $p, q, r$  distinct prime numbers.

*Proof.* Let  $G$  be a  $K_{3,3}$ -free group, and let  $N$  be a minimal normal subgroup of  $G$ . First, consider the case  $|G| = p^4q$ . Clearly,  $|N| = q$ . Since the number of the subgroups of  $G/N$  is at most 6,  $G/N$  is isomorphic to  $\mathbb{Z}_{p^4}$ . Let  $A < B < C < D$  be a chain of non-trivial  $p$ -subgroups of  $G$ . Then, one may form  $NA$ ,  $NB$  and  $NC$  which are proper subgroups. As the orders of those groups are different, they form a  $K_7$  in  $\Gamma(G)$  (intersection of any two of them contains  $A$ ). Therefore,  $G$  is not  $K_{3,3}$ -free. Similar arguments can also be applied for  $|G| = p^5q$  case.

Next, suppose  $|G| = p^2qr$ . Clearly,  $N$  is not a  $p$ -group. Without loss of generality we may assume that  $|N| = r$ . Then  $|G/N| = p^2q$ . If  $G/N$  is not cyclic, then the number of subgroups of  $G/N$  exceeds 6. This is clear if Sylow  $p$ -subgroup of  $G/N$  is elementary abelian. And if  $G/N$  is not abelian, then there must be a non-normal subgroup (since there is no subgroup of  $G/N$  isomorphic to  $Q_8$ , it is not Hamiltonian) implying there are more than 6 subgroups. Hence,  $G/N$  must be cyclic. In this case there are subgroups of  $G/N$  of orders  $p^2$ ,  $pq$ ,  $p$ , and  $q$ . Then, by the Correspondence Theorem there are four subgroups containing  $N$ , say  $A$ ,  $B$ ,  $C$ ,  $D$  of orders  $p^2r$ ,  $pqr$ ,  $pr$ , and  $qr$  respectively. Let  $T$  be a subgroup of order  $p^2q$ . By the product formula  $T$  intersects  $A$ ,  $B$ , and  $C$  non-trivially. That is,  $A, B, C, D, N, T$  span a subgraph in  $\Gamma(G)$  containing  $K_{3,3}$ .  $\square$

Now we examine the other cases.

**Lemma 4.** *Let  $G$  be a non-nilpotent group of order  $p^3q$  where  $p$  and  $q$  are distinct prime numbers. Then,  $G$  is  $K_{3,3}$ -free if and only if it is isomorphic to*

$$\mathbb{Z}_q \rtimes_{\alpha} \mathbb{Z}_{p^3} = \langle a, b \mid a^q = b^{p^3} = 1, bab^{-1} = a^{\alpha} \rangle$$

where  $p^3$  divides  $q - 1$  and  $\alpha$  is any integer not divisible by  $q$  whose order in the unit group  $\mathbb{Z}_q^*$  of  $\mathbb{Z}_q$  is  $p^3$ .

*Proof.* Suppose that  $G$  is  $K_{3,3}$ -free. Clearly, the order of the minimal normal subgroup  $N$  cannot be  $p^2$ . Therefore, we only need to consider the following two cases.

*Case I:*  $|N| = p$ . Since the number of subgroups of  $G/N$  is at most 6, we have  $G/N \cong \mathbb{Z}_{p^2q}$ . Let  $P$  be a Sylow  $p$ -subgroup of  $G$  containing  $N$ ; and let  $Q$  be a Sylow  $q$ -subgroup. By the Correspondence Theorem  $P$  is the unique Sylow  $p$ -subgroup containing  $N$ , hence it is normal in  $G$ . As  $G$  is not a nilpotent group,  $Q$  is not a normal subgroup of  $G$ . Let  $H$  and  $K$  be the subgroups containing  $N$  with orders  $pq$  and  $p^2q$  respectively. Since  $NQ^g$  is a subgroup of order  $pq$  for any conjugate  $Q^g$  of  $Q$  which contains  $N$  and since  $H$  is the unique subgroup of order  $pq$  containing  $N$  by the Correspondence Theorem,  $H$  contains all conjugates of  $Q$  and this implies the number of Sylow  $q$ -subgroups of  $G$  is  $|H : N| = p$ . In particular  $H \cong \mathbb{Z}_p \rtimes \mathbb{Z}_q$  and  $p > q$ . Then, the normalizer  $N_G(Q)$  has order  $p^2q$ . Moreover, since  $N_G(Q)$  is self-normalizing by Remark 3, there are  $p$  conjugates of  $N_G(Q)$  which are different from the normal subgroup  $K$ . By the product formula,  $P, H, K$  and  $p$  conjugates of  $N_G(Q)$  pairwise intersect non-trivially. In other words, those subgroups form a  $K_{p+3}$  on  $\Gamma(G)$ . As  $p \geq 3$ ,  $G$  cannot be  $K_{3,3}$ -free in this case.

*Case II:*  $|N| = q$ . Since  $|G : N| = p^3$  and since the number of subgroups of  $G/N$  is at most six,  $G/N$  is isomorphic to either  $\mathbb{Z}_{p^3}$  or  $Q_8$ . Notice that a group with a unique maximal subgroup is necessarily cyclic and by Remark 1 a non-cyclic  $p$ -group has at least three maximal subgroups. Therefore,  $G/N$  must have a unique minimal subgroup even if it is not cyclic.

*Case II (a):*  $G/N \cong \mathbb{Z}_{p^3}$ . Take three non-trivial  $p$ -subgroups  $A < B < C$  and form  $NA$  and  $NB$ . As the orders of those groups are different, they form a  $K_5$  in  $\Gamma(G)$ . Also, since  $G$  is not nilpotent, there are more than one Sylow  $p$ -subgroups of  $G$ . If  $A$  is contained by a Sylow  $p$ -subgroup  $D$  other than  $C$ , then together with  $D$  we have 6 proper non-trivial subgroups mutually intersecting non-trivially. On the other hand, if any two Sylow  $p$ -subgroups intersect trivially, then  $\Gamma(G)$  is  $K_{3,3}$ -free. Notice that  $NA$  is the unique subgroup of  $G$  of order  $pq$  and  $NB$  is the unique subgroup of  $G$  of order  $p^2q$ . Let  $Q = \langle a \rangle$  and  $P = \langle b \rangle$ . We want to write a presentation for  $G$ . Since  $Q$  is normal,  $bab^{-1} = a^{\alpha}$  for some integer  $\alpha$  not divisible by  $q$ . Observe that,  $b^k ab^{-k} = a^{\alpha^k}$  for any integer  $k$ . This implies  $\alpha^{p^3} \equiv 1 \pmod{q}$ , i.e. the order of  $\alpha$  in the unit group  $\mathbb{Z}_q^*$  divides  $p^3$ . Moreover, its order is exactly  $p^3$ , as otherwise, the intersection of some Sylow  $p$ -subgroups would be non-trivial. Conversely, the group

$$\mathbb{Z}_q \rtimes_{\alpha} \mathbb{Z}_{p^3} = \langle a, b \mid a^q = b^{p^3} = 1, bab^{-1} = a^{\alpha} \rangle$$

has the subgroup structure described above and it is  $K_{3,3}$ -free. See Figure 4.

*Case II (b):*  $G/N \cong Q_8$ . Then, there are 5 non-trivial subgroups of a Sylow  $p$ -subgroup each containing a unique minimal subgroup  $A$ . Together with  $NA$  we have 6 subgroups forming a  $K_6$  in  $\Gamma(G)$ . Thus, there is no  $K_{3,3}$ -free group in this case.  $\square$




Figure 4:  $\Gamma(\mathbb{Z}_q \rtimes_{\alpha} \mathbb{Z}_{p^3})$ , gray vertices represents subgroups of orders  $q$ ,  $pq$ , and  $p^2q$

There are non-nilpotent solvable planar groups of orders

$$p^2q, \quad p^2q^2, \quad pqr, \quad \text{and} \quad pq$$

which are necessarily  $K_{3,3}$ -free. Previously, we proved that the groups presented at the second and third items of the following lemma are planar.

**Lemma 5** (see [8, Lemma 4.6]). *Let  $G$  be a non-nilpotent group of order  $p^2q$  where  $p$  and  $q$  are distinct prime numbers. Then,  $G$  is  $K_{3,3}$ -free if and only if it is isomorphic to one of the following groups:*

1.

$$\mathbb{Z}_3 \rtimes \mathbb{Z}_4, \quad \text{or} \quad D_{18},$$

2.

$$\mathbb{Z}_q \rtimes_{\alpha} \mathbb{Z}_{p^2} = \langle a, b \mid a^q = b^{p^2} = 1, bab^{-1} = a^{\alpha} \rangle$$

where  $p^2$  divides  $q - 1$  and  $\alpha$  is any integer not divisible by  $q$  whose order in the unit group  $\mathbb{Z}_q^*$  of  $\mathbb{Z}_q$  is  $p^2$ ,

3.

$$(\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes_{\beta} \mathbb{Z}_q = \langle a, b, c \mid a^p = b^p = c^q = 1, ab = ba, cac^{-1} = b, cbc^{-1} = a^{-1}b^{\beta} \rangle$$

where  $q$  divides  $p + 1$  and  $\beta$  is any integer such that the matrix  $\theta = \begin{bmatrix} 0 & -1 \\ 1 & \beta \end{bmatrix}$  has order  $q$  in the group  $GL(2, \mathbb{Z}_p)$  and such that  $\theta$  has no eigenvalue in  $\mathbb{Z}_p$ .

*Proof.* Suppose that  $G$  is  $K_{3,3}$ -free. There are three possible cases for the order of the minimal normal subgroup  $N$  of  $G$ .

*Case I:*  $|N| = p$ . Let  $P$  be the Sylow  $p$ -subgroup of  $G$ , and  $Q$  be a Sylow  $q$ -subgroup of  $G$ .

*Case I (a):*  $P$  is not a normal subgroup of  $G$ . Then  $N$  is contained in every Sylow  $p$ -subgroups as well as in some subgroups of order  $pq$ . However, there are  $q = 1 + kp$  conjugates of  $P$  and since  $G$  is  $K_6$ -free, we have  $p = 2$ ,  $q = 3$  and  $H := NQ$  must be a normal subgroup of  $G$ . Notice that, the three Sylow  $p$ -subgroups together with  $N$  and  $H$  form a  $K_5$  in  $\Gamma(G)$ . Moreover,  $Q$  is a normal subgroup of  $G$ ; otherwise,  $H \cong \mathbb{Z}_p \rtimes \mathbb{Z}_q$ ,

however  $3 \nmid 2 - 1$ . If  $P \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , there would be a non-normal subgroup  $K \cong \mathbb{Z}_2$ , as otherwise,  $P$  would be a normal subgroup of  $G$ . Then  $QK$  is connected by an edge with two of the three Sylow  $p$ -subgroups as well as with  $H$  which is a contradiction because we assumed that  $G$  is  $K_{3,3}$ -free. Therefore,  $P \cong \mathbb{Z}_4$  and we can easily observe that

$$\mathbb{Z}_3 \rtimes \mathbb{Z}_4 = \langle a, b \mid a^3 = b^4 = 1, bab^{-1} = a^2 \rangle$$

is  $K_{3,3}$ -free as it has exactly six proper non-trivial subgroups and the minimal subgroup of order 3 has degree one in the intersection graph.

*Case I (b):*  $P$  is the normal Sylow  $p$ -subgroup of  $G$ . As  $G$  is not a nilpotent group by assumption,  $Q$  is not a normal subgroup of  $G$ .

Suppose that there is a normal subgroup  $L$  of  $G$  of order  $pq$  containing  $Q$ . Then  $L$  contains all conjugates of  $Q$ , hence  $L \cong \mathbb{Z}_p \rtimes \mathbb{Z}_q$  and in particular  $q \mid p - 1$ . Moreover, by Remark 3 any subgroup containing  $N_G(Q)$  is self-normalizing and this implies  $N_G(Q) \neq Q$ , as  $L \triangleleft G$  by assumption. However,  $N_G(Q) \neq G$  either, thus  $H := N_G(Q)$  is of order  $pq$  and it is not a normal subgroup of  $G$ . Let  $K$  be the subgroup of  $H$  of order  $p$ . Notice that since  $p > q$ , we have  $K \triangleleft H$ . Clearly, conjugates of  $H$  together with  $K$  and  $P$  form a  $K_{p+2}$  in  $\Gamma(G)$ . Therefore  $p = 3$  and  $q = 2$ . However, any (Sylow)  $q$ -subgroup is contained by the normal subgroup  $L$  implying there is an edge between  $L$  and any conjugate of  $H$ . That is, conjugates of  $H$  together with  $K$ ,  $P$ , and  $L$  span a subgraph containing  $K_{3,3}$ .

Now suppose that there is no normal subgroup of order  $pq$ . In particular  $NQ$  is not a normal subgroup of  $G$ . As in the previous paragraph, conjugates of  $NQ$  together with  $N$  and  $P$  form a  $K_{p+2}$  in  $\Gamma(G)$ . Therefore  $p = 3$  and  $q = 2$ , as the number of Sylow  $q$ -subgroups is  $\equiv 1 \pmod{q}$ . (Since any subgroup of index 2 must be normal,  $p \neq 2$ .) If  $P \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ , there must be a normal subgroup  $K$  of order  $p$  different from  $N$ . To see this, consider the action of  $Q$  by conjugation on the set of subgroups of order  $p$ . (Notice that there are totally four subgroups of order  $p$ .) Since  $N$  is a normal subgroup of order  $p$  and the length of an orbit of  $Q$  is either 1 or 2, there must be a subgroup  $K$  fixed by  $Q$  and different from  $N$ . However,  $G$  is generated by the elements of  $N$ ,  $K$ , and  $Q$ , thus  $K$  is a normal subgroup. Then  $KQ$  is a group of order  $pq$  different from  $NQ$  and its conjugates. This is because,  $NQ$  and  $KQ$  have unique subgroups of order  $p$  which are not conjugate to each other. By the product formula any two subgroups of order  $pq$  intersects non-trivially. Therefore, conjugates of  $NQ$  together with the conjugates of  $KQ$  form a  $K_6$  in  $\Gamma(G)$ . Finally, if  $P \cong \mathbb{Z}_9$ , we have the dihedral group

$$D_{18} = \langle a, b \mid a^9 = b^2 = 1, bab = a^{-1} \rangle$$

which is  $K_{3,3}$ -free. See Figure 5.

*Case II:*  $|N| = q$ . As the Sylow  $q$ -subgroup  $N$  is normal and as  $G$  is not a nilpotent group, there are at least three Sylow  $p$ -subgroups, say  $P_i$  ( $1 \leq i \leq q$ ).

Suppose that  $G/N \cong \mathbb{Z}_p \times \mathbb{Z}_p$ . By the Correspondence Theorem there are at least three subgroups  $H_j$  ( $1 \leq j \leq p + 1$ ) of order  $pq$  each containing  $N$ . By the product formula,  $P_i \cap H_j \neq 1$  for any  $1 \leq i, j \leq 3$  and we have six vertices which span a subgraph of  $\Gamma(G)$  containing a  $K_{3,3}$ , contradiction!

Now suppose that  $G/N \cong \mathbb{Z}_{p^2}$ . If  $X = P_i \cap P_j$  is non-trivial for some distinct Sylow  $p$ -subgroups, then  $X$  must be a normal subgroup of  $G$  as  $N_G(X)$  contains both  $P_i$  and  $P_j$ . However, this case was considered in Case I (a). If the intersection of any pair of Sylow  $p$ -subgroups is trivial, then  $G$  has a presentation

$$\mathbb{Z}_q \rtimes_{\alpha} \mathbb{Z}_{p^2} = \langle a, b \mid a^q = b^{p^2} = 1, bab^{-1} = a^{\alpha} \rangle$$

and it is planar. See [8, Lemma 4.6] for details.

*Case III:*  $|N| = p^2$ . As the Sylow  $p$ -subgroup  $N$  is normal and as  $G$  is not a nilpotent group, any subgroup of order  $q$  is not normal in  $G$ . We want to observe that there are no subgroups of  $G$  of order  $pq$ . To see this, first suppose that there is a subgroup  $H$  of  $G$  of order  $pq$ . If  $H$  is a normal subgroup of  $G$ , obviously  $H$  contains all (Sylow)  $q$ -subgroups. Then  $A = H \cap N$  is normal in  $H$  as well as in  $G$ , since  $N$  is abelian and  $\langle H, N \rangle = G$ . However, this is in contradiction with the assumption that there is no normal subgroup of order  $p$ . If  $H$  is not a normal subgroup of  $G$ , since a subgroup of smallest prime index must be normal, we have  $p > q$ . Then, again  $A$  is a normal subgroup of  $H$  and of  $G$  and we have the same contradiction. Therefore, there is no subgroup of  $G$  of order  $pq$ . In that case,  $G$  has a presentation

$$(\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes_{\beta} \mathbb{Z}_q = \langle a, b, c \mid a^p = b^p = c^q = 1, ab = ba, cac^{-1} = b, cbc^{-1} = a^{-1}b^{\beta} \rangle$$

and it is planar. See [8, Lemma 4.6] for details.  $\square$




Figure 5:  $\Gamma(D_{18})$ , gray colored vertices represents subgroups of order  $pq$

**Lemma 6** (see [8, Lemma 4.7]). *Let  $G$  be a non-nilpotent group of order  $p^2q^2$  where  $p > q$  are distinct prime numbers. Then,  $G$  is  $K_{3,3}$ -free if and only if it is isomorphic to*

$$(\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes_{\beta} \mathbb{Z}_{q^2} = \langle a, b, c \mid a^p = b^p = c^{q^2} = 1, ab = ba, cac^{-1} = b, cbc^{-1} = a^{-1}b^{\beta} \rangle$$

where  $q^2$  divides  $p + 1$  and  $\beta$  is any integer such that the matrix  $\theta = \begin{bmatrix} 0 & -1 \\ 1 & \beta \end{bmatrix}$  has order  $q^2$  in the group  $GL(2, \mathbb{Z}_p)$  and such that  $\theta^q$  has no eigenvalue in  $\mathbb{Z}_p$ .

*Proof.* Suppose that  $G$  is  $K_{3,3}$ -free. First we shall observe that the minimal normal subgroup  $N$  of  $G$  must be a Sylow subgroup. To this end, let us assume  $|N| = p$ . Then,  $G/N \cong \mathbb{Z}_{pq^2}$  and there exists a unique Sylow  $p$ -subgroup  $P$  containing  $N$ . Since Sylow  $p$ -subgroups are conjugate and since  $N$  is a normal subgroup,  $P$  is also a normal subgroup of  $G$ . Let  $Q$  be a Sylow  $q$ -subgroup. By assumption  $G$  is not nilpotent, hence  $Q$  is not a normal subgroup of  $G$ . (Notice that assuming  $|N| = q$ , one may deduce in a similar fashion that the unique Sylow  $q$ -subgroup is a normal subgroup of  $G$ . However, this is not possible for  $p > q$ .)

Let  $A, B, C$  be the subgroups of respective orders  $pq, pq^2, p^2q$  containing  $N$  and  $Q_i$  ( $1 \leq i \leq 3$ ) be three Sylow  $q$ -subgroups. Let  $X$  be a group of order  $q$ . Since  $N$  is a normal subgroup,  $NX$  is a group of order  $pq$  containing  $X$ . This implies  $A$  contains any

group of order  $q$ , as it is the unique subgroup of order  $pq$  containing  $N$ . That is  $A \cap Q_i$  is non-trivial for  $1 \leq i \leq 3$ . This is also true for  $B$  and  $C$ , and so  $A, B, C$  together with  $Q_i$  span a subgraph containing a  $K_{3,3}$  in  $\Gamma(G)$ .

By the preceding discussion we conclude that  $N$  is the normal Sylow  $p$ -subgroup of  $G$  and since it is minimal,  $N \cong \mathbb{Z}_p \times \mathbb{Z}_p$  by Remark 4. We know that if the rank of the minimal normal subgroup  $N$  is two, then  $G/N$  has at most three subgroups, hence it is a cyclic group of prime or prime squared order. Since the order of  $G$  is  $p^2q^2$ , we conclude  $Q \cong \mathbb{Z}_{q^2}$  where  $Q$  is a Sylow  $q$ -subgroup. Let  $K$  be the unique subgroup of  $G$  of order  $p^2q$  containing  $N$ . Since any subgroup of order  $p^2q$  contains  $N$ , we see that  $K$  is the unique subgroup of  $G$  of order  $p^2q$ . Moreover, since  $NX = K$  for any subgroup  $X$  of order  $q$ ,  $K$  contains all subgroups of  $G$  of order  $q$ . Also, since any subgroup of a  $K_{3,3}$ -free group is also  $K_{3,3}$ -free,  $K$  is isomorphic to the third group stated in the previous Lemma 5. In particular, there is no subgroup of  $K$  of order  $pq$  and in turn this implies there are no subgroups of  $G$  of order  $pq^2$  or  $pq$ . To see this observe that if  $H < G$  is of order  $pq$ , then  $H \cap K$  contains a subgroup of order  $p$  by the product formula. However,  $K$  contains every subgroup of order  $q$  which implies  $H < K$ , contradiction! Similar argument works when  $|H| = pq^2$ . Hence  $G$  is a group with a normal Sylow  $p$ -subgroup isomorphic to  $\mathbb{Z}_p \times \mathbb{Z}_p$  and a non-normal Sylow  $q$ -subgroup isomorphic to  $\mathbb{Z}_{q^2}$  and there are no subgroups of  $G$  of order  $pq$  or of order  $pq^2$ . Such a group has a presentation

$$(\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes_{\beta} \mathbb{Z}_{q^2} = \langle a, b, c \mid a^p = b^p = c^{q^2} = 1, ab = ba, cac^{-1} = b, cbc^{-1} = a^{-1}b^{\beta} \rangle$$

and it is planar. See [8, Lemma 4.7] for details.  $\square$

**Lemma 7** (see [8, Lemma 4.8]). *Let  $G$  be a non-nilpotent group of order  $pqr$  where  $p < q < r$  are distinct prime numbers. Then,  $G$  is  $K_{3,3}$ -free if and only if it is isomorphic to*

$$\mathbb{Z}_r \rtimes_{\alpha} \mathbb{Z}_{pq} = \langle a, b \mid a^r = b^{pq} = 1, bab^{-1} = a^{\alpha} \rangle$$

where  $pq$  divides  $r - 1$  and  $\alpha$  is any integer not divisible by  $r$  whose order in the unit group  $\mathbb{Z}_r^*$  of  $\mathbb{Z}_r$  is  $pq$ .

*Proof.* Let  $R$  be a Sylow  $r$ -subgroup. Applying Sylow theorems it can be easily observed that  $R \triangleleft G$ . Since  $|G/R| = pq$  and  $p < q$ , we see that  $G/R$  is either abelian (hence cyclic) or isomorphic to  $\mathbb{Z}_q \rtimes \mathbb{Z}_p$ . In the latter case the number of subgroups of  $G/R$  is  $q+3$ . Since this number must be at most six,  $G$  is isomorphic to the symmetric group  $S_3 \cong \mathbb{Z}_3 \rtimes \mathbb{Z}_2$  in the latter case.

Suppose that  $G/R \cong S_3$ . By the Correspondence Theorem, there is a unique subgroup  $N$  of order  $3r$  and three subgroups  $K_i$  ( $1 \leq i \leq 3$ ) of order  $2r$  containing  $R$ . Since  $N$  is a Hall  $\{3, r\}$ -subgroup and  $R$  is normal,  $N$  is a normal subgroup of  $G$  as well (see Remark 5). Let  $Q$  be the Sylow 3-subgroup of  $G$  contained by  $N$ . Then, by Remark 3,  $Q$  is a normal subgroup of  $G$ . Let  $H$  be a Hall  $\{2, 3\}$ -subgroup of  $G$ . If  $H$  is not a normal subgroup of  $G$ , then the number of its conjugates is  $|G : H| = r$  and those subgroups together with  $Q$  form a  $K_{r+1}$  in  $\Gamma(G)$ . Since  $r \geq 5$ , the intersection graph cannot be  $K_{3,3}$ -free in this case. Also, if  $H \triangleleft G$  then it is easy to observe that the subgroups  $H$ ,  $N$ ,  $R$  together with  $K_i$  ( $1 \leq i \leq 3$ ) form a subgraph containing  $K_{3,3}$  in the intersection graph.

Suppose that  $G/R \cong \mathbb{Z}_{pq}$ . By the Correspondence Theorem, there are unique subgroups  $N$  of order  $pr$  and  $M$  of order  $qr$ . As in the preceding paragraph both  $M$  and  $N$  are normal subgroups. Let  $K$  be a subgroup of order  $pq$ . Clearly,  $K$  is not a normal subgroup of  $G$  and in particular it has  $r$  conjugates. Now assume that there exist two

distinct conjugates  $K_1$  and  $K_2$  of  $K$  such that their intersection  $X = K_1 \cap K_2$  is non-trivial. Then, as  $|X|$  is either  $p$  or  $q$ , we have  $X \triangleleft G$ ; and this implies  $X$  is contained by all conjugates of  $K$ . That is, conjugates of  $K$  together with  $X$  form a  $K_{r+1}$  in the intersection graph which is a contradiction as  $r \geq 5$ . Therefore, any two distinct subgroup of order  $pq$  intersects trivially. Such a group has a presentation

$$\mathbb{Z}_r \rtimes_{\alpha} \mathbb{Z}_{pq} = \langle a, b \mid a^r = b^{pq} = 1, bab^{-1} = a^{\alpha} \rangle$$

and it is planar. See [8, Lemma 4.8] for details.  $\square$

Finally, intersection graph of any group of order  $pq$  consists of isolated vertices and so  $K_{3,3}$ -free. For further references we state it as a lemma.

**Lemma 8.** *Let  $G$  be a non-nilpotent group of order  $pq$  where  $p$  and  $q$  are distinct prime numbers and  $p > q$ . Then,  $q \mid p - 1$  and*

$$G \cong \mathbb{Z}_p \rtimes \mathbb{Z}_q$$

*is  $K_{3,3}$ -free.*

## Non-solvable groups

First, we shall show that there is no finite non-abelian simple group which is  $K_{3,3}$ -free. To this end, we need the following result.

**Theorem 9** ([6, Theorem 1]). *If the finite group  $G$  contains a maximal subgroup  $M$  which is nilpotent of class less than 3, then  $G$  is solvable.*

As a consequence of Theorem 9, if a Sylow  $p$ -subgroup  $P$  of  $G$  is maximal and  $|P| = p^3$ , then  $G$  is solvable.

**Proposition 10.** *If  $G$  is a finite minimal simple group, then  $\Gamma(G)$  contains a  $K_{3,3}$  as a subgraph.*

*Proof.* Consider a finite simple group  $U$  which is  $K_{3,3}$ -free. Then there exists a minimal finite simple group  $G$  which is isomorphic to a non-abelian composition factor of some subgroup of  $U$ . Thus,  $G$  must be  $K_{3,3}$ -free.

Minimal simple groups are known (see [4, Corollary 1]). Thus,  $G$  is isomorphic to one of the following groups:  $PSL_2(q)$ ,  $Sz(q)$ ,  $PSL_3(3)$ .

In view of Feit-Thompson Theorem, 2 divides  $|G|$ . Let  $S$  be a Sylow 2-subgroup of  $G$ . Then  $S$  is a 2-group from Lemmas 1 or 2. Thus, either  $S \cong \mathbb{Z}_{2^i}$ , where  $1 \leq i \leq 6$ , or  $S \in \{\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_4 \times \mathbb{Z}_2, D_8, Q_8\}$ .

Since the intersection graphs of each of the 2-groups  $\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $D_8$  and  $Q_8$  contains  $K_{1,3}$ , those groups must be maximal in  $G$ . By Theorem 9,  $G$  is solvable. A contradiction.

Suppose that  $S$  is cyclic. Then, by Remark 6,  $S$  has a normal complement in  $G$  which contradicts with the assumption that  $G$  is simple.

Thus,  $S \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ . By the Burnside normal complement theorem (see [10, Theorem 7.50]), the normalizer  $N_G(S)$  properly contains  $S$ . Clearly,  $N_G(S)$  is a proper subgroup of  $G$ , as otherwise,  $S$  would be a normal subgroup.

Normalizers of Sylow 2-subgroups of finite simple groups are known (see [3, Corollary]). Thus,  $G$  is isomorphic to either  $PSL_2(q)$ , where  $q \equiv \pm 3 \pmod{8}$  (in this case  $N_G(S) \cong A_4$ ), or to  $PSL_3(3)$ . But  $PSL_3(3)$  properly contains  $S_4$  (see [2]), therefore is not  $K_{3,3}$ -free. If  $G \cong PSL_2(q)$ , where  $q \equiv \pm 3 \pmod{8}$ , then there is a subgroup  $H \cong D_{q \pm 1}$  of  $G$  which is a subgroup of odd index (see [1, Table 8.7]). Take  $S$  to be a Sylow 2-subgroup of  $H$ . Then  $H$ ,  $N_G(S)$ ,  $S$  and three proper subgroups of  $S$  form a graph which contains  $K_{3,3}$ . A contradiction.  $\square$

**Corollary 11.** *A finite non-solvable group is not  $K_{3,3}$ -free.*

*Proof.* Let  $G$  be a finite non-solvable group. Since  $G$  has a non-abelian simple composition factor which is not  $K_{3,3}$ -free by Proposition 10,  $G$  is not  $K_{3,3}$ -free as well.  $\square$

Our main result follows from Lemmas 1, 2, 4, 5, 6, 7, 8 and Corollary 11.

## References

- [1] Bray J.N., Holt D.F., Roney-Dougal C.M., *The maximal subgroups of the low-dimensional finite classical groups*, Cambridge: Cambridge Univ. Press, 2013.
- [2] Conway J. H., Curtis R. T., Norton S. P., Parker R. A., Wilson R. A., *Atlas of finite groups*, Oxford: Clarendon Press, 1985
- [3] Kondrat'ev A. S., *Normalizers of the Sylow 2-Subgroups in Finite Simple Groups*, Mathematical Notes. 2005. Vol. 78, no. 3. P. 338-346
- [4] Thompson J. G., *Nonsolvable finite groups all of whose local subgroups are solvable*, Bull. Amer. Math. Soc. 1968. Vol. 74, no. 3. P. 383-437
- [5] Berkovich Y., *Groups of prime power order. Vol. 1*, de Gruyter Expositions in Mathematics, vol. 46, Walter de Gruyter GmbH & Co. KG, Berlin, 2008, With a foreword by Zvonimir Janko.
- [6] Deskins W. E., *A condition for the solvability of a finite group*, Illinois J. Math. **5** (1961), 306–313.
- [7] Gorenstein D., *Finite groups*, second ed., Chelsea Publishing Co., New York, 1980.
- [8] Kayacan S. and Yaraneri E., *Finite groups whose intersection graphs are planar*, J. Korean Math. Soc. **52** (2015), no. 1, 81–96.
- [9] Rajkumar R. and Devi P., *Toroidality and projective-planarity of intersection graphs of subgroups of finite groups*, ArXiv e-prints (2015).
- [10] Rotman, J. J., *An introduction to the theory of groups*, fourth ed., Graduate Texts in Mathematics, vol. 148, Springer-Verlag, New York, 1995.